

# *Salmonella enterica* serotypes isolated from squabs reveal multidrug resistance and a distinct pathogenicity gene repertoire

K.M. Osman <sup>(1)\*</sup>, S.H. Marouf <sup>(1)</sup>, O.A. Mehana <sup>(2)</sup> & N. AlAtfeehy <sup>(3)</sup>

(1) Department of Microbiology, Faculty of Veterinary Medicine, Cairo University, P.O. Box 12211, Cairo, Egypt

(2) Biotechnology Unit, Animal Research Institute, Ministry of Agriculture, 7 Nadi Al Said Street, Dokki, Egypt

(3) Department of Poultry Diseases, Animal Research Institute, Ministry of Agriculture, 7 Nadi Al Said Street, Dokki, Egypt

\*Corresponding author: kamelia-osman@hotmail.com

## Summary

The consumption of squab (young unfledged pigeons) as part of the cuisine of many countries, together with the observation that squabs are vectors of zoonotic agents, may make them a public health risk. This study was designed to determine the serotypes, distribution of 11 virulence genes (*invA*, *avrA*, *ssaQ*, *mgcC*, *siiD*, *sopB*, *gipA*, *sodC1*, *sopE1*, *spvC*, *bcfC*) and the antimicrobial resistance profiles of salmonellae recovered from squabs. Six isolates were identified from among 45 (13.3%) squabs sampled. Three serotypes were identified according to the Kauffmann–White serotyping scheme: *Salmonella* Typhimurium (4/6; 66.7%), *S. Braenderup* (1/6; 16.7%) and *S. Lomita* (1/6; 16.7%). Polymerase chain reaction analyses revealed the presence of *invA*, *sopB* and *bcfC* in all six isolates, whereas *sopE1* and *gipA* were absent. All six isolates were resistant to lincomycin and streptomycin, but all were susceptible to ciprofloxacin, colistin sulphate and gentamicin. Among the *S. Typhimurium* isolates, seven resistance profiles were identified: penicillins, aminoglycosides, fluoroquinolones, lincosamides, phenicols, tetracyclines and sulphonamides; four resistance profiles were identified in the isolates of *S. Braenderup* and *S. Lomita*: aminoglycosides, fluoroquinolones, lincosamides and polymyxin. Thus, the distribution of resistance to the antibiotics was largely dependent on serotype identity. The presence of *invA*, *avrA*, *ssaQ*, *mgcC*, *siiD*, *sopB* and *bcfC* was associated with resistance to chloramphenicol; *invA*, *sopB* and *bcfC* with resistance to streptomycin and lincosamide; and *invA* and *sodC1* with resistance to trimethoprim–sulfamethoxazole. The identification of serotypes *S. Typhimurium*, *S. Braenderup* and *S. Lomita* in the squab samples has important implications because these serotypes are significant causes of food poisoning and enteric fever in humans.

## Keywords

Antibiotic resistance – Pigeon – *Salmonella* Braenderup – *Salmonella* Lomita – *Salmonella* Typhimurium – Squab – Virulence gene.

## Introduction

The first historical mention of domesticated pigeons being used for food was in Egypt in 3,000 BC (1). Since their initial domestication, pigeons have been regarded as a cheap source of good meat, particularly the squab meat, which is valued highly. The Romans also kept pigeons for food and were familiar with the practice of force-feeding squabs in order to fatten the young birds more quickly. Wild pigeons

were hunted as a cheap and readily available source of protein, although the meat from older and wild pigeons is tougher and requires a long period of stewing or roasting to tenderise. Usually considered a delicacy, squab is tender, moist and richer in taste than many commonly consumed poultry meats. Today, squab is part of the cuisine of many countries, including France, the United States (USA), Italy, Northern Africa, and several Asian countries (1, 2, 3). The Egyptians, Chinese and other Asians consider squab meat an aphrodisiac. In the USA, squab is 'increasingly a specialty

item', as the larger and cheaper chicken has displaced it (4). Nevertheless, squab produced from specially raised utility pigeons continues to grace the menus of American haute cuisine restaurants such as Le Cirque and the French Laundry (5, 6) and has enjoyed endorsements from some celebrity chefs (1). Accordingly, squab is often sold for much higher prices than other poultry, sometimes as high as US\$8 per pound weight (1).

Human salmonellosis has been associated with contaminated poultry and with direct contact with food animals, highlighting the possibility of transmission of these organisms to humans through the food chain (7). Although there are few reports of disease transmission from pigeons to humans (8), their close interaction, together with the observation that these birds are vectors for zoonotic agents (9), may make them a public health risk. Squab meat is regarded as safer than some other poultry products, as it harbours fewer pathogens (2, 10), but there has not yet been a detailed study of the role of squab as a vector of *Salmonella* species.

To determine the extent to which squabs might harbour salmonellae and pose a risk to the human population in Egypt, squabs on sale in Cairo markets were screened for the presence of salmonellae relevant to public health. Isolates were examined for the presence of 11 virulence-associated genes (*invA*, *avrA*, *ssaQ*, *mgtC*, *siiD*, *sopB*, *gipA*, *sodC1*, *spvC*, *sopE1*, *bcfC*) and their antimicrobial resistance profiles were determined.

## Methods

### Sampling and isolation

The research was carried out in the Department of Poultry Diseases, Animal Research Institute, Dokki, Egypt (30° 3' North, 31° 15' East). The birds used in the study comprised 45 clinically healthy squabs randomly selected and purchased from local markets in Cairo and delivered alive to the laboratory. Sterile culettes (Product No. 4360210, Becton, Dickinson & Co., Cockeysville, Maryland, USA) were used to collect cloacal swabs for faecal culture. All swabs were immediately transported to the laboratory in ice-cold containers and processed within 6 h of collection. Strains of salmonellae were isolated according to the standard ISO-6579 method (11).

### Serotyping of salmonellae

Purified isolates were serotyped using a slide agglutination method with O- and H-group antigen-specific antisera (Denka-Seiken, Tokyo, Japan). Once the antigenic formulae were obtained, the Kauffmann–White serotyping scheme was used to identify the serotype.

### Antimicrobial susceptibility testing using the disc diffusion method

The antimicrobial resistance profiles of the *S. Typhimurium*, *S. Braenderup* and *S. Lomita* isolates were determined using discs impregnated with a range of antibiotics at fixed concentrations (Becton, Dickinson & Co.). The following antimicrobials were chosen on the basis of their common use in treating and preventing *Salmonella* spp. infection in poultry and humans, and because they represent a range of antimicrobial classes: ampicillin (10 mg), amoxicillin (20 mg), gentamicin (10 mg), neomycin (30 µg), streptomycin (10 mg), lincomycin (30 µg), chloramphenicol (30 mg), colistin (10 mg), tetracycline (30 mg), trimethoprim (5 µg), sulfamethoxazole + trimethoprim (23.75 mg + 1.75 mg). In addition, resistance was determined against three broad-spectrum fluoroquinolones: ciprofloxacin (5 µg), norfloxacin (5 µg) and nalidixic acid (30 mg).

Each isolate was inoculated onto a Muller–Hinton agar (Oxoid) plate and antibiotic discs were placed on the surface of the agar (six discs per plate). The plates were incubated for 24 h to 48 h at 37°C. After incubation, the diameter of the halos was measured to assess resistance or susceptibility according to the interpretation criteria for *Escherichia coli* (*E. coli*) (ATCC No. 25922) established by the Clinical Laboratory Standards Institute (12). Multidrug resistance is defined as resistance to two or more antibiotics belonging to different antibiotic classes (9).

### Detection of virulence determinants

A polymerase chain reaction (PCR) amplification assay (for primers and assay conditions, see Table I) was used to detect virulence genes in all the isolates. Briefly, isolates were grown overnight at 37°C in Luria–Bertani broth (Sigma, St. Louis, Missouri, USA). Genomic DNA was extracted using a boiling method and used as the target for PCR assays. Lysates were chilled on ice and then spun for 5 min in a microcentrifuge at 14,000 g to pellet the debris. Portions (5 µl) of the supernatants were used as templates in the PCR. The isolates were screened for the presence of 11 virulence genes (*invA*, *avrA*, *ssaQ*, *mgtC*, *siiD*, *sopB*, *gipA*, *sodC1*, *sopE1*, *spvC*, *bcfC*). The PCR conditions and primers used for the target genes have been previously published (13, 14, 15). The specificity of the oligonucleotide primers was confirmed using strains of *S. Enteritidis* (ATCC 13076) and *E. coli* (ATCC 50034) as reference samples.

### Statistical analysis

The distribution of resistance phenotypes among the serotypes, either resistance to a single antibiotic or multidrug resistance, was tested using a contingency table analysis where rows = serotype, and columns = resistance versus susceptibility. The significance of homogeneity in antimicrobial resistance patterns among

**Table I**  
**Virulence factor targets and primers, including nucleotide sequences, polymerase chain reaction conditions and references**

| Gene designation    | Location on SPI and gene function   | Oligonucleotide sequences (5'-3')   | PCR conditions <sup>(a)</sup>   |                  |                                 | Product size (bp) | Reference |
|---------------------|---|---|---------------------------------|------------------|---------------------------------|-------------------|-----------|
|                     |   |   | Denaturing                      | Annealing        | Extension                       |                   |           |
| <i>invA</i>         | Type III secretion system apparatus SPI-1: Invasion of macrophages  | gtg aaa tta tcg cca cgt tcg<br>ggc aa<br>tca tcg cac cgt caa agg<br>aac g | 72°C for<br>30 s <sup>(b)</sup> | 64°C for<br>30 s | 94°C for<br>60 s                | 284               | 13        |
| <i>avrA</i>         | SPI-1: Controls <i>Salmonella</i> -induced inflammation   | cct gta ttg ttg agc gtc ttg<br>aga aga gct tcg ttg aat<br>gtc c           | 95°C for<br>30 s                | 58°C for<br>30 s | 72°C for<br>30 s <sup>(b)</sup> | 422               | 14        |
| <i>ssaQ</i>         | SPI-2: Secretion system apparatus protein, component of second T3SS   | gaa tag cga atg aag agc<br>gtc gtc c<br>cat cgt gtt atc ctc tgt<br>cag c  | "                               | "                | "                               | 455               | 14        |
| <i>mgtC</i>         | SPI-4: Mg <sup>2+</sup> uptake  | tga cta tca atg ctc cag<br>tga at<br>att tac tgg ccg cta tgc<br>tgt tg    | "                               | "                | "                               | 677               | 14        |
| <i>siiD (Spi4D)</i> | Type I secretion: SPI-4   | gaa tag aag aca aag cga<br>tca tc<br>gct ttg ttc acg cct ttc atc          | "                               | "                | "                               | 655               | 15        |
| <i>sopB</i>         | SPI-5: Inositol polyphosphate phosphatase that promotes macropinocytosis, regulates SCV localisation and promotes fluid secretion | tca gaa gRc gtc taa cca<br>ctc<br>tac cgt cct cat gca cac tc              | "                               | "                | "                               | 517               | 14        |
| <i>gipA</i>         | Gifsy-1 bacteriophage: Peyer's patch-specific virulence factor  | acg act gag cag cgt gag<br>ttg gaa atg gtg acg gta<br>gac                 | "                               | "                | "                               | 518               | 14        |
| <i>sodC1</i>        | Gifsy-2 bacteriophage: Periplasmic Cu-, Zn-superoxide dismutases  | cgg gca gtg ttg aca aat<br>aaag<br>tgt tgg aat tgt gga gtc                | "                               | "                | "                               | 424               | 14        |
| <i>sopE1</i>        | Cryptic bacteriophage: Promotes membrane ruffling and disrupts tight junctions  | act cct tgc aca acc aaa<br>tgc gga<br>tgt ctt ctg cat ttc gcc acc         | "                               | "                | "                               | 422               | 14        |
| <i>spvC</i>         | pSLT: A phosphothreonine lyase required for complete virulence in murine models   | acc aga gac att gcc ttc c<br>ttc tga tcg ccg cta ttc g                    | "                               | "                | "                               | 467               | 14        |
| <i>bcfC</i>         | Chromosome: Bovine colonisation factor, fimbrial usher  | acc aga gac att gcc ttc c<br>ttc tgc tcg ccg cta ttc g                    | 72°C for<br>30 s <sup>(b)</sup> | 53°C for<br>30 s | 95°C for<br>30 s                | 467               | 14        |

a) Polymerase chain reaction, 35 cycles

b) After 30 cycles, final extension step of 4 min at 72°C

bp: base pairs

PCR: polymerase chain reaction

SCV: *Salmonella*-containing vacuole

SPI: *Salmonella* pathogenicity islands

T3SS: Type III secretion system

the groups was assessed using Pearson's  $\chi^2$  exact test and SAS version 9.2 software (SAS, Cary, New Jersey, USA). Findings were considered significant where  $p < 0.05$ . A high  $\chi^2$  value indicates that the resistant phenotypes are not proportionately distributed among serotypes, some serotypes having a higher frequency of resistant phenotypes than others. To assess the significance of differences (D) between individual D values, 95% confidence intervals (95% CIs) were calculated. The normal approximation to the binomial distribution was used to calculate the 95% CIs for the proportion of isolates resistant to individual antimicrobial agents.

## Results

A total of six (13.3%) isolates of salmonellae were identified from the 45 squabs. Serotyping of the isolates differentiated them into *S. Typhimurium* (4/6; 66.7%), *S. Braenderup* (1/6; 16.7%) and *S. Lomita* (1/6; 16.7%).

### Virulotyping

All isolates were screened using PCR analysis for the presence or absence of 11 selected virulence genes (Table II). The genes *invA* and *sopB* (carried by *Salmonella* pathogenicity islands [SPIs]) and *bcfC* (fimbria-related) were detected in all six isolates. In contrast, *sopE1* (encoding a translocated effector protein) and *gipA* (encoding a Peyer's patch-specific virulence factor) were absent. The gene *spvC*, carried by the *Salmonella* virulence plasmid, was present in three of the four isolates of *S. Typhimurium* but was absent from both the Braenderup and Lomita isolates. The Braenderup and Lomita isolates carried *avrA* (located in SPI-1) and *mgcC* (carried by SPIs), both of which were absent from the Typhimurium isolates. In contrast, *sodC1* (located on a bacteriophage) was identified in the Typhimurium isolates only.

### Combinations of virulence genes

Over all, four different combinations of virulence genes were detected among the serotypes (Table III). The four Typhimurium isolates possessed three combinations of virulence genes, two of them being identical (*sodC1*, *sopB*, *spvC* and *bcfC*).

### Distribution of resistance to individual antimicrobial agents

Frequent resistance to the antimicrobials tested was evident in all six isolates of salmonellae (Table IV). In particular, all isolates were resistant to lincomycin and streptomycin but all were susceptible to ciprofloxacin and gentamicin; resistance patterns to the other antimicrobials exhibited great diversity. The four Typhimurium isolates showed seven

different resistance profiles (penicillin, aminoglycosides, fluoroquinolones, lincosamides, phenicols, tetracyclines, sulphonamides). Interestingly, four of the resistance profiles of Braenderup and Lomita isolates were the same as for Typhimurium (aminoglycosides, fluoroquinolones, lincosamides, polymyxin). Significant correlations among various antimicrobials at isolate level are shown in Tables III and IV. The distribution of resistance to the antibiotics was largely dependent on serotype identity. Resistance to colistin sulphate ( $\chi^2 = 6.0$ ;  $p < 0.014$ ) was associated with serotypes Braenderup and Lomita.

### Association of antimicrobial resistance phenotype with virulence-associated genes

Analysis of the presence of the 11 virulence-associated genes in the tested isolates is shown in Table III. Detailed analysis revealed associations of resistance/susceptibility phenotypes with potential virulence genes. The association of genes *invA*, *sopB* and *bcfC* with particular antimicrobial resistance phenotypes (streptomycin and lincosamide) has not been recorded previously. The presence of *invA* and *sodC1* was associated with resistance to trimethoprim-sulfamethoxazole.

## Discussion

Although many bacteria in poultry and poultry-related samples have been identified, to the authors' knowledge and from a thorough scrutiny of the literature, no studies have yet reported on virulence genes associated with different *Salmonella* serotypes and/or antimicrobial resistance in salmonellae recovered from squabs.

Notably, serotypes *S. Braenderup* and *S. Lomita* have never been detected in squabs until now. The age of the bird does not influence the *Salmonella* infection rate in pigeons and squabs (10); however, having a parent that tests positive for salmonellae greatly increases the odds of testing positive as a squab, which may imply transovarian transmission or transmission through intimate contact, either by the faecal-oral route or possibly from the crop of the parent to the oral cavity of the squab (10). The successful isolation of serotypes Typhimurium, Braenderup and Lomita reported here has important implications, because they are a significant cause of food poisoning and enteric fever in humans (16), which makes the spread of Braenderup and Lomita among pigeons a concern for human health. Both these serotypes have been isolated from cases of food poisoning in England and Wales (17), Germany (18), Switzerland (19), the USA (20) and Japan (21, 22). With regard to isolates derived from squabs, rodents acting as reservoirs for the bacteria help to maintain and spread the infection in the pigeon population; for example, by contaminating food sources.

**Table II**  
**Distribution of virulence genes among three *Salmonella* serotypes**

| <i>Salmonella</i> serotypes | Virulence genes |             |             |             |             |             |             |              |              |             |             |
|-----------------------------|-----------------|-------------|-------------|-------------|-------------|-------------|-------------|--------------|--------------|-------------|-------------|
|                             | <i>invA</i>     | <i>avrA</i> | <i>ssaQ</i> | <i>mgtC</i> | <i>siiD</i> | <i>sopB</i> | <i>gipA</i> | <i>sodC1</i> | <i>sopE1</i> | <i>spvC</i> | <i>bcfC</i> |
| Typhimurium                 | +               | ND          | ND          | ND          | ND          | +           | ND          | +            | ND           | ND          | +           |
| Typhimurium                 | +               | ND          | +           | ND          | +           | +           | ND          | +            | ND           | +           | +           |
| Typhimurium                 | +               | ND          | ND          | ND          | ND          | +           | ND          | +            | ND           | +           | +           |
| Typhimurium                 | +               | ND          | ND          | ND          | ND          | +           | ND          | +            | ND           | +           | +           |
| Braenderup                  | +               | +           | +           | +           | +           | +           | ND          | ND           | ND           | ND          | +           |
| Lomita                      | +               | +           | +           | +           | +           | +           | ND          | ND           | ND           | ND          | +           |

ND: not detected

**Table III**  
**Distribution of virulence gene combinations and antibiotic resistance phenotypes in three *Salmonella* serotypes**

| Serotype    | Virulence gene combination <sup>(a)</sup>   | Resistance phenotype                                 |
|-------------|---|--|
| Typhimurium | <b><i>invA</i>, <i>sodC1</i>, <i>sopB</i>, <i>bcfC</i></b>  | Amo, Amp, Chl, Lin, Str, Na, Neo, Nor, Tet, Tri, Sxt |
|             | <b><i>invA</i>, <i>ssaQ</i>, <i>siiD</i>, <i>sodC1</i>, <i>sopB</i>, <i>spvC</i>, <i>bcfC</i></b> | Lin, Str, Tet, Sxt                                   |
|             | <b><i>invA</i>, <i>sodC1</i>, <i>sopB</i>, <i>spvC</i>, <i>bcfC</i></b>                           | Amo, Neo, Chl, Lin, Str, Na, Tet, Sxt                |
|             | <b><i>invA</i>, <i>sodC1</i>, <i>sopB</i>, <i>spvC</i>, <i>bcfC</i></b>                           | Chl, Lin, Str, Na, Tet, Sxt                          |
| Braenderup  | <b><i>invA</i>, <i>avrA</i>, <i>ssaQ</i>, <i>mgtC</i>, <i>siiD</i>, <i>sopB</i>, <i>bcfC</i></b>  | Chl, Lin, Str, Na, Tet                               |
| Lomita      | <b><i>invA</i>, <i>avrA</i>, <i>ssaQ</i>, <i>mgtC</i>, <i>siiD</i>, <i>sopB</i>, <i>bcfC</i></b>  | Chl, Lin, Str, Na                                    |

a) Bold indicates consistent frequency of virulence genes  
 Amo: amoxicillin  
 Amp: ampicillin  
 Chl: chloramphenicol

Lin: lincomycin  
 Na: nalidixic acid  
 Neo: neomycin  
 Nor: norfloxacin

Str: streptomycin  
 Tet: tetracycline  
 Tri: trimethoprim  
 Sxt: trimethoprim-sulfamethoxazole

Many of the virulence factors in salmonellae are present as part of pathogenicity islands on the chromosome, although some virulence factors are encoded on plasmids or bacteriophages (23). In the present study, the *Salmonella* invasion gene *invA*, which is essential for entry of the bacteria into epithelial cells, is a putative inner membrane component of the SPI-1-dependent Type III secretion system (TTSS-1) virulence apparatus (24) and was present in all six isolates. The virulotyping results for isolates carrying the *Salmonella* genomic island-1 (SGI-1) indicated that little or no variation occurred for most genes incorporated in SPIs and for the fimbrial marker. Thus, the genes *sopB* and *bcfC* were also present throughout all serotypes. It was observed that some virulence patterns were serotype specific; for example, the isolates of Braenderup and Lomita shared a common virulotype that was characterised by the presence of seven genes and the absence of *sodC1* and *spvC*. The association of a certain serotype with a virulence-associated gene panel was also confirmed by the microarray results reported by Beutlich *et al.* (25) and a previous study by Huehn *et al.* (14) of European isolates belonging to different serotypes of salmonellae.

All the isolates of Typhimurium, Braenderup and Lomita that were tested showed multidrug resistance. Nevertheless,

all isolates were sensitive to gentamicin and ciprofloxacin, and sensitivity to ampicillin, amoxicillin, neomycin, norfloxacin, chloramphenicol, tetracycline, trimethoprim and trimethoprim-sulfamethoxazole was also frequent. Ciprofloxacin is the drug of choice for treating potentially life-threatening infections caused by multidrug-resistant strains of salmonellae in adult humans (26). However, recent studies have documented the emergence of ciprofloxacin resistance in salmonellae isolated from foods (27, 28), with nalidixic acid resistance indicating the first step in the development of resistance to ciprofloxacin (29). Thus, nalidixic-acid-resistant serotypes of Typhimurium, Braenderup and Lomita usually show decreased sensitivity to ciprofloxacin (29, 30). In the present study, however, although the isolates were resistant to nalidixic acid they remained sensitive to ciprofloxacin. For comparison, it is noteworthy that isolates from pigeon eggs were sensitive to ciprofloxacin, norfloxacin, gentamicin and chloramphenicol (100%), but resistant to tetracycline (50%) (31). Realistically, antibiotic resistance can be considered a virulence factor (32). Resistance mechanisms are pandemic and create an enormous clinical and financial burden on healthcare systems worldwide (32). The almost complete absence of resistance to ciprofloxacin and norfloxacin among the isolates tested is important, because these two

**Table IV**  
**Distribution of resistance to individual antimicrobial agents among three *Salmonella enterica* serotypes isolated from squabs**

| Antimicrobials                 | Distribution of resistance to antimicrobials |               |              |                                |               |              |                            |               |              |
|--------------------------------|--|---------------|--------------|--------------------------------|---------------|--------------|----------------------------|---------------|--------------|
|                                | <i>S. Typhimurium</i> (n = 4/6)              |               |              | <i>S. Braenderup</i> (n = 1/6) |               |              | <i>S. Lomita</i> (n = 1/6) |               |              |
|                                | No. of resistant isolates                    | % of isolates | 95% CI       | No. of resistant isolates      | % of isolates | 95% CI       | No. of resistant isolates  | % of isolates | 95% CI       |
| <b>Penicillins</b>             |  |               |              |                                |               |              |                            |               |              |
| Ampicillin                     | 2/4  | 50            | 9.18, 90.81  | 0/1                            | 0             | 10.78, 94.54 | 0/1                        | 0             | 10.78, 94.54 |
| Amoxicillin                    | 2/4  | 50            | 9.18, 90.81  | 0/1                            | 0             | 10.78, 94.54 | 0/1                        | 0             | 10.78, 94.54 |
| <b>Aminoglycosides</b>         |  |               |              |                                |               |              |                            |               |              |
| Gentamicin                     | 0/4  | 0             | 2.35, 60.42  | 0/1                            | 0             | 10.78, 94.54 | 0/1                        | 0             | 10.78, 94.54 |
| Neomycin                       | 1/4  | 25            | 1.32, 78.06  | 0/1                            | 0             | 10.78, 94.54 | 0/1                        | 0             |              |
| Streptomycin                   | 4/4  | 100           | 39.58, 97.65 | 1/1                            | 100           | 5.46, 89.22  | 1/1                        | 100           | 5.46, 89.22  |
| <b>Fluoroquinolones</b>        |  |               |              |                                |               |              |                            |               |              |
| Ciprofloxacin                  | 0/4  | 0             | 2.35, 60.42  | 0/1                            | 0             | 10.78, 94.54 | 0/1                        | 0             |              |
| Nalidixic acid                 | 3/4  | 75            | 21.94, 98.68 | 1/1                            | 100           | 5.46, 89.22  | 1/1                        | 100           | 5.46, 89.22  |
| Norfloxacin                    | 1/4  | 25            | 1.32, 78.06  | 0/1                            | 0             | 10.78, 94.54 | 0/1                        | 0             |              |
| <b>Lincosamides</b>            |  |               |              |                                |               |              |                            |               |              |
| Lincomycin                     | 4/4  | 100           | 39.58, 97.65 | 1/1                            | 100           | 5.46, 89.22  | 1/1                        | 100           | 5.46, 89.22  |
| <b>Phenicol</b>                |  |               |              |                                |               |              |                            |               |              |
| Chloramphenicol                | 3/4  | 75            | 21.94, 98.68 | 0/1                            | 0             | 10.78, 94.54 | 0/1                        | 0             | 10.78, 94.54 |
| <b>Polymyxin</b>               |  |               |              |                                |               |              |                            |               |              |
| Colistin sulphate              | 0/4  | 0             | 2.35, 60.42  | 1/1                            | 100           | 5.46, 89.22  | 1/1                        | 100           | 5.46, 89.22  |
| <b>Tetracyclines</b>           |  |               |              |                                |               |              |                            |               |              |
| Tetracycline                   | 3/4  | 75            | 21.94, 98.68 | 0/1                            | 0             | 10.78, 94.54 | 0/1                        | 0             | 10.78, 94.54 |
| <b>Sulphonamides</b>           |  |               |              |                                |               |              |                            |               |              |
| Trimethoprim                   | 2/4  | 50            | 9.18, 90.81  | 0/1                            | 0             | 10.78, 94.54 | 0/1                        | 0             | 10.78, 94.54 |
| Trimethoprim-sulfa-methoxazole | 3/4  | 75            | 21.94, 98.68 | 0/1                            | 0             | 10.78, 94.54 | 0/1                        | 0             | 10.78, 94.54 |

CI: 95% binomial confidence interval (95% CI) based on normal approximation

antimicrobials are the primary agents used against invasive salmonellosis cases in humans (33).

In Senegal and Gambia, significant associations have been found between the presence of some virulence genes and resistance to commonly used antimicrobials (34). The isolates of Typhimurium, Braenderup and Lomita in the present investigation have a core pattern of resistance to the four antimicrobials lincomycin, streptomycin, nalidixic acid and colistin sulphate, referred to as quattuor-resistance type LSNaCs. The five genes for this resistance type, *invA*, *ssaQ*, *siiD*, *sopB* and *bcfC*, are clustered within the genome on the genomic island SGI-1, which also contains horizontally transferable genetic elements such as phage- and plasmid-related genes (35, 36).

An inevitable side effect of antimicrobial use is the emergence and dissemination of resistance in pathogenic bacteria, with the possibility of resistant zoonotic bacteria reaching the intestinal tract of humans (37, 38). The emergence and development of resistance to lincomycin, streptomycin, nalidixic acid and colistin sulphate has been linked to the wide use of antimicrobials in veterinary practice (39, 40, 41, 42, 43) for the treatment and prevention of disease and for growth promotion.

## Conclusion

Understanding the associations between human cases of salmonellosis, animal sources and the environment is

epidemiologically important in attempts to control the spread of these infections within communities (7, 44). The isolation of *S. enterica* serotypes Typhimurium, Braenderup and Lomita from squabs has important implications, because they are significant causes of food poisoning and enteric fever in humans (16, 17, 18, 19, 20, 21, 22). Carriage of

the virulence-associated genes *invA*, *sopB*, and *bcfC* among multidrug-resistant strains may increase the propensity of such strains to be of major clinical relevance, particularly if specific virulence factors are also carried (33).

## Les sérotypes de *Salmonella enterica* isolés à partir de pigeonneaux témoignent de l'apparition de multi-résistances et d'un répertoire spécifique de gènes codant pour la pathogénicité

K.M. Osman, S.H. Marouf, O.A. Mehana & N. AlAtfeehy

### Résumé

Le pigeonneau est une denrée de choix dans la culture gastronomique de nombreux pays ; il est aussi le vecteur de plusieurs maladies zoonotiques et représente donc un facteur potentiel de risque pour la santé publique. Les auteurs présentent les résultats d'une étude visant à déterminer les sérotypes des salmonelles isolées à partir de pigeonneaux, la distribution au sein de ces sérotypes de 11 gènes de virulence (*invA*, *avrA*, *ssaQ*, *mgtC*, *siiD*, *sopB*, *gipA*, *sodC1*, *sopE1*, *spvC*, *bcfC*) et les profils de résistance aux agents antimicrobiens. Six isolats ont été obtenus à partir des échantillons prélevés sur 45 pigeonneaux (13,3 %). Trois sérotypes ont été identifiés en utilisant la méthode de typage basée sur le schéma de Kauffmann-White : *Salmonella* Typhimurium (4/6 ; 66,7 %) ; *S. Braenderup* (1/6 ; 16,7 %) et *S. Lomita* (1/6 ; 16,7 %). La technique d'amplification en chaîne par polymérase a révélé que chacun des six isolats était pourvu des gènes *invA*, *sopB* et *bcfC* mais dépourvu des gènes *sopE1* et *gipA*. Les six isolats étaient résistants à la lincomycine et à la streptomycine, mais sensibles à la ciprofloxacine, au sulfate de colistine et à la gentamicine. Sept profils de résistance ont été identifiés parmi les *S. Typhimurium* isolées : pénicillines, aminoglycosides, fluoroquinolones, lincosamides, phénicols, tétracyclines et sulphonamides ; quatre profils de résistance ont été identifiés dans les souches isolées de *S. Braenderup* et *S. Lomita* : aminoglycosides, fluoroquinolones, lincosamides et polymyxines. Ainsi, la distribution des résistances aux antibiotiques dépendait fortement du sérotype. Les corrélations suivantes ont été constatées : présence des gènes *invA*, *avrA*, *ssaQ*, *mgtC*, *siiD*, *sopB* et *bcfC* et résistance au chloramphénicol ; présence des gènes *invA*, *sopB* et *bcfC* et résistance à la streptomycine et à la lincosamide ; présence des gènes *invA* et *sodC1* et résistance au triméthoprime-sulfaméthoxazole. L'identification des sérotypes *S. Typhimurium*, *S. Braenderup* et *S. Lomita* dans les prélèvements issus de pigeonneaux est une observation importante car ces sérotypes sont responsables de toxi-infections alimentaires et de fièvres typhoïdes chez l'homme.

### Mots-clés

Gène de la virulence – Pigeon – Pigeonneau – Résistance aux antibiotiques – *Salmonella* Braenderup – *Salmonella* Lomita – *Salmonella* Typhimurium.

## Serotipos de *Salmonella enterica* aislados en pichones revelan multirresistencia a los medicamentos y un repertorio característico de genes de patogenicidad

K.M. Osman, S.H. Marouf, O.A. Mehana & N. AlAtfeehy

### Resumen

Su consumo como parte de la gastronomía de muchos países, así como la observación de que son vectores de agentes zoonóticos, pueden hacer de los pichones (crías implumes de paloma) un riesgo para la salud pública. Los autores describen un estudio en el que se determinaron los serotipos, la distribución de 11 genes de virulencia (*invA*, *avrA*, *ssaQ*, *mgtC*, *siiD*, *sopB*, *gipA*, *sodC1*, *sopE1*, *spvC* y *bcfC*) y las características de antibiorresistencia de salmonelas aisladas a partir de seis (un 13,3%) de los 45 pichones analizados. Aplicando el método de clasificación de Kauffmann–White se caracterizaron tres serotipos: *Salmonella* Typhimurium (4/6; 66,7%), *S. Braenderup* (1/6; 16,7%) y *S. Lomita* (1/6; 16,7%). Los análisis por reacción en cadena de la polimerasa (PCR) revelaron la presencia de los genes *invA*, *sopB* y *bcfC* en las seis bacterias aisladas, que carecían en cambio de *sopE1* y *gipA*. Los seis microorganismos eran resistentes a la lincomicina y la estreptomina y sensibles al ciprofloxacino, el sulfato de colistina y la gentamicina. Entre los microorganismos *S. Typhimurium* aislados se observaron siete perfiles de resistencia (a las penicilinas, los aminoglicósidos, las fluoroquinolonas, las lincosamidas, los fenicoles, las tetraciclinas y las sulfonamidas), y en *S. Braenderup* y *S. Lomita* se observaron cuatro (aminoglicósidos, fluoroquinolonas, lincosamidas y polimixina). La distribución de las antibiorresistencias dependía mucho, por consiguiente, del serotipo de que se tratara. La presencia de los genes *invA*, *avrA*, *ssaQ*, *mgtC*, *siiD*, *sopB* y *bcfC* se acompañaba de resistencia al cloranfenicol; la de *invA*, *sopB* y *bcfC*, de resistencia a la estreptomina y la lincosamida; y la de *invA* y *sodC1*, de resistencia al trimetoprim–sulfametoxazol. El descubrimiento de los serotipos *S. Typhimurium*, *S. Braenderup* y *S. Lomita* en las muestras de pichones tiene importantes consecuencias, porque estos serotipos son una causa sustancial de intoxicaciones alimentarias y fiebres tifoideas en el ser humano.

### Palabras clave

Antibiorresistencia – Gen de virulencia – Paloma – Pichón – *Salmonella* Braenderup – *Salmonella* Lomita – *Salmonella* Typhimurium.

## References

- Blechman A.D. (2006). – Pigeons: the fascinating saga of the world's most revered and reviled bird. Open City Books, Grove Press, New York, 256 pp.
- Morgan J.L. (2006). – Culinary creation: an introduction to foodservice and world cuisine. Butterworth-Heinemann hospitality management series. Butterworth-Heinemann, Oxford, 220 pp.
- Natala A.J., Asemadahun N.D., Okubanjo O.O., Ulayi B.M., Owolabi Y.H., Jato I.D. & Yusuf K.H. (2009). – A survey of parasites of domesticated pigeon (*Columba livia* domestic) in Zaria, Nigeria. *Int. J. soft Comput.*, 4 (4), 148–150.
- Keller T. (1999). – The French laundry cookbook. Artisan, New York, 325 pp.

5. Colin J. (2007). – Animal archeology: domestic pigeons and the nature-culture dialectic. *Qual. sociol. Rev.*, **3** (1), 74–95.
6. Bruni F. (2008). – In defense of decadence. *New York Times*, 6 February 2008. Available at: [events.nytimes.com/2008/02/06/dining/reviews/06rest.html?scp=1&sq=squab&st=nyt](http://events.nytimes.com/2008/02/06/dining/reviews/06rest.html?scp=1&sq=squab&st=nyt) (accessed on 24 February 2008).
7. Khaitsa M.L. & Doetkott D. (2012). – Antimicrobial drug resistance and molecular characterization of *Salmonella* isolated from domestic animals, humans and meat products; *Salmonella* – a dangerous foodborne pathogen, Chapter 11. In *Immunology and microbiology* (S.M. Barakat Mahmoud, ed.). InTech, Rijeka, Croatia, 450 pp. doi:10.5772/1308.
8. Haag-Wackernagel D. & Moch H. (2004). – Health hazards posed by feral pigeons. *J. Infect.*, **48**, 307–313.
9. Hoelzer K., Soyer Y., Rodriguez-Rivera L.D., Cummings K.J., McDonough P.L., Schoonmaker-Bopp G.J., Root T.P., Dumas N.B., Warnick L.D., Grohn Y.T., Wiedmann M., Baker K.N.K., Besser T.E., Hancock D.D. & Davis M.A. (2010). – The prevalence of multidrug resistance is higher among bovine than human *Salmonella enterica* serotype Newport, Typhimurium, and 4,5,12:i:- isolates in the United States but differs by serotype and geographic region. *Appl. environ. Microbiol.*, **76**, 5947–5959.
10. Jeffrey J.S., Atwill E.R. & Hunter A. (2001). – Farm and management variables linked to fecal shedding of *Campylobacter* and *Salmonella* in commercial squab production. *Poult. Sci.*, **80**, 66–70.
11. Osman K.M., Yousef A.M.M., Aly M.M. & Radwan M.I. (2010). – *Salmonella* spp. infection in imported 1-day-old chicks, ducklings, and turkey poults: a public health risk. *Foodborne Path. Dis.*, **7**, 383–390.
12. Clinical Laboratory Standards Institute (CLSI) (2008). – Performance standards for antimicrobial susceptibility testing, 9th informational supplement. CLSI, Wayne, Pennsylvania.
13. Salehi T.Z., Mahzounieh M. & Saeedzadeh A. (2005). – Detection of *InvA* gene in isolated *Salmonella* from broilers by PCR method. *Inter. J. Poultry Sci.*, **4**, 557–559.
14. Huehn S., La Ragione R.M., Anjum M., Saunders M., Woodward M.J., Bunge C., Helmuth R., Hauser E., Guerra B., Beutlich J., Brisabois A., Peters T., Svensson L., Madajczak G., Litrup E., Imre A., Herrera-Leon S., Mevius D., Newell D.G. & Malorny B. (2010). – Virulotyping and antimicrobial resistance typing of *Salmonella enterica* serovars relevant to human health in Europe. *Foodborne Path. Dis.*, **7**, 523–535.
15. Hauser E., Hebner F., Tietze E., Helmuth R., Junker E., Prager R., Schroeter A., Rabsch W., Fruth A. & Malorny B. (2011). – Diversity of *Salmonella enterica* serovar Derby isolated from pig, pork and humans in Germany. *Int. J. Food Microbiol.*, **151**, 141–149.
16. World Health Organization (WHO) (2000). – World Health Organization Surveillance Programme for Control of Foodborne Infections and Intoxications in Europe, 7th Report 2000. WHO, Geneva.
17. Health Protection Agency (2000). – General outbreaks of foodborne illness in humans, England and Wales: quarterly report 2000. *CDR Weekly*, **498** (10), 434–444.
18. Rossier P., Urfer E., Burnens A., Bille J., Francioli P., Mean F. & Zwahlen A. (2000). – Clinical features and analysis of the duration of colonization during an outbreak of *Salmonella* Braenderup gastroenteritis. *Schweizer. Med. Wochenschr.*, **130**, 1185–1191.
19. Urfer E., Rossier P., Mean F., Krending M., Burnens A., Bille J., Francioli P. & Zwahlen A. (2000). – Outbreak of *Salmonella* Braenderup gastroenteritis due to contaminated meat pies: clinical and molecular epidemiology. *Clin. Microbiol. Infect.*, **6**, 536–542.
20. Gupta S., Nalluswami K., Snider C., Perch M., Balasegaram M., Bumeister D., Lockett J., Sandt C., Hoekstra R. & Montgomery S. (2007). – Outbreak of *Salmonella* Braenderup infections associated with Roma tomatoes, northeastern United States, 2004: a useful method for subtyping exposures in field investigations. *Epidemiol. Infect.*, **135**, 1165–1173.
21. Anon. (2009). – Salmonellosis in Japan as of June 2009. *Infect. Agents Surveill. Rep.*, **30** (8).
22. Mizoguchi Y., Suzuki E., Tsuchida H., Tsuda T., Yamamoto E., Nakase E. & Doi H. (2011). – Outbreak of *Salmonella* Braenderup infection originating in boxed lunches in Japan in 2008. *Acta Med. Okayama*, **65**, 63–69.
23. Silva C., Wiesner M. & Calva E. (2012). – The importance of mobile genetic elements in the evolution of *Salmonella*: pathogenesis, antibiotic resistance and host adaptation. In *Salmonella* – a diversified superbug (Y. Kumar, ed.). InTech, Rijeka, Croatia. doi:10.5772/2471
24. Hur J., Choi Y.Y., Park J.H., Jeon B.W., Lee H.S., Kim A.R. & Lee J.H. (2011). – Antimicrobial resistance, virulence-associated genes, and pulsed-field gel electrophoresis profiles of *Salmonella enterica* subsp. *enterica* serovar Typhimurium isolated from piglets with diarrhea in Korea. *Can. J. vet. Res.*, **75**, 49–56.
25. Beutlich J., Jahn S., Malorny B., Hauser E., Hühn S., Schroeter A., Rodicio M.R., Appel B., Threlfall J., Mevius D., Helmuth R. & Guerra B. (2011). – Antimicrobial resistance and virulence determinants in European *Salmonella* genomic island 1-positive *Salmonella enterica* isolates from different origins. *Appl. environ. Microbiol.*, **77**, 5655–5664.
26. Hsueh P., Teng L., Tseng S., Chang C., Wan J., Yan J., Lee C., Chuang Y., Huang W., Yang D., Shyr J., Yu K., Wang L., Lu J., Ko W., Wu J., Chang F., Yang Y., Lau Y., Liu Y., Liu C., Ho S. & Luh K. (2004). – Ciprofloxacin-resistant *Salmonella enterica* Typhimurium and Choleraesuis from pigs to humans, Taiwan. *Emerg. infect. Dis.*, **10**, 60–68.

27. Mayrhofer S., Peter P., Smulders F.J.M. & Hilbert F. (2004). – Antimicrobial resistance profile of five major food-borne pathogens isolated from beef, pork and poultry. *Int. J. Food Microbiol.*, **97**, 23–29.
28. Olah P.A., Sherwood J.S., Elijah L.M., Dockter M.R., Doetkott C. & Miller Z. (2004). – Comparison of antimicrobial resistance in *Salmonella* and *Campylobacter* isolated from turkeys in the Midwest USA. *Food Microbiol.*, **21**, 779–789.
29. Van Looveren M., Chasseur-Libotte M.L., Godard C., Lammens C., Wijdooghe M., Peeters L. & Goossens H. (2001). – Antimicrobial susceptibility of nontyphoidal *Salmonella* isolated from humans in Belgium. *Acta clin. Belg.*, **56**, 180–186.
30. Wybot I., Wildemauwe C., Godard C., Bertrand S. & Collard J.M. (2004). – Antimicrobial drug resistance in nontyphoid human *Salmonella* in Belgium: trends for the period 2000–2002. *Acta clin. Belg.*, **59**, 152–160.
31. Jahantigh M. & Nili H. (2010). – Drug resistance of *Salmonella* spp. isolated from pigeon eggs. *Comp. clin. Pathol.*, **19**, 437–439.
32. Davies J. & Davies D. (2010). – Origins and evolution of antibiotic resistance. *Microbiol. mol. Biol. Rev.*, **74**, 417–433.
33. Gebreyes W.A., Thakur S., Dorr P., Tadesse D.A., Post K. & Wolf L. (2009). – Occurrence of *spvA* virulence gene and clinical significance for multidrug-resistant *Salmonella* strains. *J. clin. Microbiol.*, **47**, 777–780.
34. Dione M.M., Ikumapayi U., Saha D., Mohammed N.I., Adegbola R.A., Geerts S., Ieven M. & Antonio M. (2011). – Antimicrobial resistance and virulence genes of non-typhoidal *Salmonella* isolates in the Gambia and Senegal. *J. Infect. dev. Ctries*, **5**, 765–775.
35. Boyd D., Peters G.A., Cloeckart A., Boumedine K.S., Chaslus-Dancla E., Imberechts H. & Mulvey M.R. (2001). – Complete nucleotide sequence of a 43-kilobase genomic island associated with the multidrug resistance region of *Salmonella enterica* serovar Typhimurium DT104 and its identification in phage type DT120 and serovar Agona. *J. Bacteriol.*, **183**, 5725–5732.
36. Briggs C.E. & Fratamico P.M. (1999). – Molecular characterization of an antibiotic resistance gene cluster of *Salmonella* Typhimurium DT104. *Antimicrob. Agents Chemother.*, **43**, 846–849.
37. O'Brien T.F. (2002). – Emergence, spread, and environmental effect of antimicrobial resistance: how use of an antimicrobial anywhere can increase resistance to any antimicrobial anywhere else. *Clin. infect. Dis.*, **34**, S78–S84.
38. Collignon P., Powers J.H., Chiller T.M., Kane A.A. & Aarestrup F.M. (2009). – World Health Organization ranking of antimicrobials according to their importance in human medicine: a critical step for developing risk management strategies for the use of antimicrobials in food production animals. *Clin. infect. Dis.*, **49**, 132–141.
39. Levy S.B., FitzGerald G.B. & Maccone A.B. (1976). – Changes in intestinal flora of farm personnel after introduction of a tetracycline-supplemented feed on a farm. *N. Engl. J. Med.*, **295**, 583–588.
40. Holmberg S.D., Osterholm M.T., Senger K.A. & Cohen M.L. (1984). – Drug-resistant *Salmonella* from animals fed antimicrobials. *N. Engl. J. Med.*, **311**, 617–622.
41. Threlfall E.L., Ward L.R., Frost J.A. & Willshaw G.A. (2000). – The emergence and spread of antibiotic resistance in food-borne bacteria. *Int. J. Food Microbiol.*, **43**, 625–635.
42. McEwen S.A. & Fedorka-Cray P.J. (2002). – Antimicrobial use and resistance in animals. *Clin. infect. Dis.*, **34**, S93–S106.
43. Gebreyes W.A., Davies P.R., Turkson P.K., Morrow W.E., Funk J.A., Altier C. & Thakur S. (2004). – Characterization of antimicrobial-resistant phenotypes and genotypes among *Salmonella enterica* recovered from pigs on farms, from transport trucks, and from pigs after slaughter. *J. Food Protec.*, **67**, 698–705.
44. Hoelzer K., Switt A.I.M. & Wiedmann M. (2011). – Animal contact as a source of human non-typhoidal salmonellosis. *Vet. Rec.*, **42**, 34.